



## COVER SHEET

---

**This is the author version of article published as:**

Byrne, Nuala M. and Weinsier, Roland L. and Hunter, Gary R. and Desmond, Renee and Patterson, Mindy A. and Darnell, Betty E. and Zuckerman, Paul A. (2003) Influence of distribution of lean body mass on resting metabolic rate after weight loss and weight regain: Comparison of responses in white and black women. *The American Journal of Clinical Nutrition* 77(6):pp. 1368-1373.

**Copyright 2003 The American Society for Clinical Nutrition**

Accessed from <http://eprints.qut.edu.au>

# Influence of distribution of lean body mass on resting metabolic rate after weight loss and weight regain: comparison of responses in white and black women<sup>1,2,3,4</sup>

Nuala M Byrne, Roland L Weinsier, Gary R Hunter, Renee Desmond, Mindy A Patterson, Betty E Darnell and Paul A Zuckerman

American Journal of Clinical Nutrition, Vol. 77, No. 6, 1368-1373, June 2003  
© 2003 [American Society for Clinical Nutrition](#)

## ABSTRACT

**Background:** Little is known about the effect of weight change on regional lean body mass (LBM) distribution or on racial differences in resting metabolic rate (RMR).

**Objective:** The study compared total and regional LBM patterns in white and black women after weight loss and regain and assessed the influence of regional LBM on variances in RMR.

**Design:** Eighteen white and 22 black women who did not differ in age, weight, and height were studied 3 times: in the overweight state, after weight reduction to the normal-weight state, and after 1 y without intervention. Total and regional lean and fat masses were assessed by dual-energy X-ray absorptiometry.

**Results:** White and black women did not differ significantly in mean ( $\pm$  SD) weight loss ( $13.4 \pm 3.6$  and  $12.7 \pm 3.2$  kg, respectively) and regain ( $6.1 \pm 5.5$  and  $6.4 \pm 5.4$  kg, respectively). Black subjects had significantly less trunk LBM and significantly more limb LBM at each time point ( $P < 0.05$ ). In both races, weight regain was associated with significant increases in limb LBM ( $P < 0.05$ ) but not in trunk LBM ( $P = 0.21$ ). RMR, adjusted for total LBM and fat mass, was significantly higher in white women after weight loss ( $P < 0.01$ ) and regain ( $P < 0.01$ ). However, no racial difference was found when RMR was adjusted for LBM distribution.

**Conclusions:** In both races, trunk LBM decreased with weight loss and remained lower, despite significant weight regain, which potentially reflected decreased organ mass. Regional LBM distribution explained the racial difference in RMR.

**Key Words:** Overweight • obesity • weight loss • body composition • resting metabolic rate • lean body mass • fat mass • regional distribution • African American women

## REFERENCES

1. World Health Organization. Obesity: preventing and managing the global epidemic. Geneva, Switzerland: World Health Organization, 1998.
2. National Heart, Lung, and Blood Institute, National Institutes of Health. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults. Washington, DC: Government Printing Office, 1998.
3. Franz MJ. Managing obesity in patients with comorbidities. *J Am Diet Assoc* 1998;98:S39–43.
4. Williamson DF, Serdula MK, Anda RF, Levy A, Byers T. Current weight loss attempts in adults: goal, duration, and rate of weight loss. *Am J Public Health* 1992;82:1251–7.
5. Kuczmarski RJ, Flegal KM, Campbell SM, Johnson CL. Increasing prevalence of overweight among US adults: the National Health and Nutrition Examination Surveys, 1960 to 1991. *JAMA* 1994;272:205–11.
6. Kumanyika SK, Obarzanek E, Stevens VJ, Hebert PR, Whelton PK. Weight-loss experience of black and white participants in NHLBI sponsored clinical trials. *Am J Clin Nutr* 1991;53(suppl):1631S–8S.
7. Darga LL, Holden JH, Olson SM, Lucas CP. Comparison of cardiovascular risk factors in obese blacks and whites. *Obes Res* 1994;2:239–45.
8. Yanovski SZ, Gormally JF, Lesser MS, Gwirtsman HE, Yanovski JA. Binge eating disorder affects outcome of comprehensive very low-calorie diet treatment. *Obes Res* 1994;2:205–12.
9. Sugerman HJ, Londrey GL, Kellum JM. Weight loss with vertical banded gastroplasty and Roux-Y gastric bypass with selective vs. random assignment. *Am J Surg* 1989;157:93–102.
10. Foster GD, Wadden TA, Swain RM, Anderson DA, Vogt RA. Changes in resting energy expenditure after weight loss in obese African American and white women. *Am J Clin Nutr* 1999;69:13–7.
11. Foster GD, Wadden TA, Vogt RA. Resting energy expenditure in obese African American and Caucasian women. *Obes Res* 1997;5:1–8.
12. Albu J, Shur M, Curi M, Murphy L, Heymsfield SB, Pi-Sunyer FX. Resting metabolic rate in obese, premenopausal black women. *Am J Clin Nutr* 1997;66:531–8.

13. Jakicic JM, Wing RR. Differences in resting energy expenditure in African-American versus Caucasian overweight females. *Int J Obes Relat Metab Disord* 1998;22:236–42.
14. Gerace L, Aliprantis A, Russell M, et al. Skeletal differences between black and white men and their relevance to body composition estimates. *Am J Hum Biol* 1994;6:255–62.
15. Malina RM. Regional body composition: age, sex, and ethnic variation. In: Roche AF, Heymsfield SB, Lohman TG, eds. *Human body composition*. Champaign, IL: Human Kinetics, 1996:217–55.
16. Aloia JF, McGowan DM, Vaswani AN, Ross P, Cohn SH. Relationship of menopause to skeletal muscle mass. *Am J Clin Nutr* 1991;53:1378–83.
17. Gallagher D, Visser M, De Meersman RE, et al. Appendicular skeletal muscle mass: effects of age, gender, and ethnicity. *J Appl Physiol* 1997;83:229–39.
18. Brozek J, Grande F. Body composition and basal metabolism in man: correlation analysis versus physiological approach. *Hum Biol* 1955;27:22–31.
19. Elia M. The inter-organ flux of substrates in fed and fasted man, as indicated by atrio-venous balance studies. *Nutr Res Rev* 1991;4:3–31.
20. Elia M. Organ and tissue contribution to metabolic rate. In: Kinney JM, Tucker HN, eds. *Energy metabolism: tissue determinants and cellular corollaries*. New York: Raven, 1992:61–80.
21. Garby L, Lammert O. Between-subjects variation in energy expenditure: estimation of the effect of variation in organ size. *Eur J Clin Nutr* 1994;48:376–8.
22. Weinsier RL, Schutz Y, Bracco D. Reexamination of the relationship of resting metabolic rate to fat-free mass and to the metabolically active components of fat-free mass in humans. *Am J Clin Nutr* 1992;55:790–4.
23. Sparti A, DeLany JP, de la Bretonne JA, Sander GE, Bray GA. Relationship between resting metabolic rate and the composition of the fat-free mass. *Metabolism* 1997;46:1225–30.
24. Hunter GR, Weinsier RL, Darnell BE, Zuckerman PA, Goran MI. Racial differences in energy expenditure and aerobic fitness in premenopausal women. *Am J Clin Nutr* 2000;71:500–6.
25. Heymsfield SB, Smith R, Aulet M, et al. Appendicular skeletal muscle mass: measurement by dual-photon absorptiometry. *Am J Clin Nutr* 1990;52:214–8.

26. Jebb SA, Goldberg GR, Elia M. DEXA measurements of fat and bone mineral in relation to depth and adiposity. In: Ellis KJ, Eastman JD, eds. Human body composition: in vivo methods, models and assessment. New York: Plenum, 1993:115–9.
27. Wang Z, Visser M, Ma R, et al. Skeletal muscle mass: evaluation of neutron activation and dual-energy X-ray absorptiometry methods. *J Appl Physiol* 1996;80:824–31.
28. Aloia JF, Vaswani A, Mikhail M, Flaster ER. Body composition by dual-energy X-ray absorptiometry in black compared with white women. *Osteoporosis Int* 1999;10:114–9.
29. Aloia JF, Vaswani A, Ma R, Flaster E. Comparison of body composition in black and white premenopausal women. *J Lab Clin Med* 1997;129:294–9.
30. Lara-Castro C, Weinsier RL, Hunter GR, Desmond R. Visceral adipose tissue in premenopausal black and white women: longitudinal study of the effects of fat gain, time, and race. *Obes Res* 2002;10:868–74.
31. Weinsier RL, Hunter GR, Desmond RA, Byrne NM, Zuckerman PA, Darnell BE. Free-living activity energy expenditure in women who are successful and unsuccessful at maintaining a normal body weight. *Am J Clin Nutr* 2002;75:499–504.
32. Nelson KM, Weinsier RL, Long CL, Schutz Y. Prediction of energy expenditure from fat-free mass and fat mass. *Am J Clin Nutr* 1992;56:848–56.
33. Weir JB. New methods for calculating metabolic rate with special reference to protein metabolism. *J Physiol* 1949;109:1–9.
34. Forbes GB. Lean body mass-body fat interrelationships in humans. *Nutr Rev* 1987;45:225–31.
35. DeLany JP, Bray GA, Harsha DW, Volaufova J. Energy expenditure in preadolescent African American and white boys and girls: the Baton Rouge Children's Study. *Am J Clin Nutr* 2002;75:705–13.
36. Gannon B, DiPietro L, Poehlman ET. Do African Americans have lower energy expenditure than Caucasians? *Int J Obes Relat Metab Disord* 2000;24:4–13.
37. Hunter GR, Weinsier RL, Gower BA, Wetzstein C. Age-related decrease in resting energy expenditure in sedentary white women: effects of regional differences in lean and fat mass. *Am J Clin Nutr* 2001;73:333–7.
38. Wadden TA, Foster GD, Stunkard AJ, Conill AM. Effects of weight cycling on the resting energy expenditure and body composition of obese women. *Am J Physiol* 1996;270:E363–6.

39. Van der Kooy K, Leenen R, Seidell JC, Deurenberg P, Hautvast JG. Effect of a weight cycle on visceral fat accumulation. *Am J Clin Nutr* 1993;58:853–7.
40. Janssen I, Ross R. Effects of sex on the change in visceral, subcutaneous adipose tissue and skeletal muscle in response to weight loss. *Int J Obes Relat Metab Disord* 1999;23:1035–46.